



Exopag 10
Astrometry
for exoplanet detection




Eduardo Bendek (NASA Ames)


Why astrometry from space?




Why astrometry from space?



Astrometry and direct Imaging





Astrometry and Period






Astrometry + Direct imaging



4 Year mission,
1 Month Cadence
Astrometry only
Guyon et al, ApJ 2013.





4 Year mission,
2 Month Cadence
**Astrometry +
Coronagraphy**
Guyon et al, ApJ2013.



Astrometry + Radial Velocity



- Expands the exploration envelope, complements RV
- Solves inclination ambiguity




ESPRESSO Online – 2016
RV 10cm/s⁻¹
Earth-like planets @ aCen HZ

HIPPARCOS ASTROMETRY CONSTRAINS THE TRUE COMPANION MASS OF RADIAL VELOCITY COMPANIONS




From Johannes Sahlmann, ESA

Sahlmann et al., 2011, A&A, 525



Radial velocities:
orbit inclination unconstrained
Brown dwarf ?



RV + Hipparcos astrometry:
orbit inclination determined
M-dwarf !

Astrometry and exoplanets...




- Great science:
 - Measure planet masses
 - Solve system inclination (AND can see any inclination)
 - Highly efficient when used with Direct Imaging
 - Complementary with RV discovery space
 - Better suited for longer period planets (closer to HZ)
- $1\mu\text{as} < \text{required for earth-like measurements}$
- $10\mu\text{as}$ enables super-earths and Neptunes
- Sparse field capability required

European astrometry missions




Hipparcos – ESA 1989 - 1993




- 0.001 μas for 117,000 stars
- 0.03 as for 2.5 million stars (Tycho2)
- 2.5 million stars
- 300Ly range

GAIA ESA 2013 - 2018



- **8 μas per visit for stars $6 < m_v < 12$**
- **25 μas per visit for stars $m_v = 15$**
- 70 visits in 5 years.
- 1000 million stars
- 30.000Ly range
- Up to 7 thousand exoplanets to be detected

Astrometry missions



10 -100 μas astrometry required to access statistical samples of exoplanets

Earth twin detection requires 0.5-1 μas

HST Astrometry




From Adam Reiss

Wide field

- WFC3/UVIS Point and stare mode $\sim 400\mu\text{as}$
- HST FGS $\sim 300\mu\text{as}$

Narrow field


- Precision Astrometry with Spatial Scanning $\sim 25\mu\text{as}$



Limiting factor




From Michael Perryman



schematic of a distorted reference frame

...it has proven impossible to eliminate these local distortions from small field observations (photographic plate or CCD), even using the method of 'block adjustment' (Eichhorn 1988)





Hipparcos - ground-based (FK5)
systematic errors (Schwan 2002)

Distortion and Astrometry



- How distortions affect astrometry
 - Cause local plate scale changes
 - Bias the astrometric measurements
- Diffractive spikes can calibrate the plate scale





NEAT



From Malbet F., Leger A., Shao M., et al.


- Remove non-pupil optics: 2 spacecrafts, 1m off-axis aperture
- Add interferometric calibration for detectors and pixels.



Mission name	Mirror diameter (m)	Focal length (m)	Field of view diameter (deg)	Focal Plane size (cm)	Ref. star mean magnitude (R mag)	DMA in 1h (μ as)	# targets for a given mass limit		
							$0.5M_{\oplus}$	$1 M_{\oplus}$	$5 M_{\oplus}$
NEAT plus	1.2	50	0.45	40	11.5	0.7	7	100	200
NEAT	1.0	40	0.56	40	11	0.8	5	70	200
NEAT light	0.8	30	0.71	35	10.5	1.0	4	50	200
EXAM	0.6	20	0.85	30	10.1	1.4	2	35	200

DMA = Differential astrometric Measurement Accuracy (rms)

Distortion and Astrometry




Courtesy: NASA

Diffraction Pupil Concept



- Concept and anticipated performance

- Astrometry accuracy: $\sim 0.3 \mu\text{as}$ (quadratic with FoV and Aperture)
Guyon et al., ApJ 200:11, 2012





Expected Single Measurement Astrometric Accuracy as a Function of Telescope Diameter and Field of View (Galactic Pole Pointing, 2 Day Integration)

Tel. Diam.	Field of View			
	0.03 deg ² ($\phi = 0.2$ deg)	0.1 deg ² ($\phi = 0.36$ deg)	0.3 deg ² ($\phi = 0.6$ deg)	1 deg ² ($\phi = 1.13$ deg)
1.0 m	0.99 μas	0.54 μas	0.31 μas	0.17 μas
1.4 m	0.62 μas	0.34 μas	0.20 μas	0.11 μas
2.0 m	0.38 μas	0.21 μas	0.12 μas	0.066 μas
2.8 m	0.24 μas	0.13 μas	0.076 μas	0.041 μas
4.0 m	0.15 μas	0.081 μas	0.047 μas	0.026 μas
5.7 m	0.092 μas	0.050 μas	0.029 μas	0.016 μas
8.0 m	0.059 μas	0.032 μas	0.019 μas	0.010 μas

- For a $D=2.4\text{m}$ and $\text{FoV}=0.03 \text{ deg}^2$
 $0.32 \mu\text{as}$ (Vis)
- Assuming:
 - 2 Day integration
 - Galactic pole pointing
 - Telescope roll
 - Diffraction limited PSF

First Laboratory Test



Relevant work or proposal	Astrometric accuracy		Start TRL	End TRL
	λ/D	Equivalent accuracy for $D=2.4\text{m}$ and $\lambda=500\text{nm}$		
First lab demonstration	3.42×10^{-3}	$147\mu\text{as}^*$	1	3
This TDEM Milestone	2.35×10^{-4}	$10\mu\text{as}^*$	3	4
Mission goal	2.35×10^{-5}	$1\mu\text{as}$	>7	9




* The lab test for these cases uses $D=0.016\text{m}$.

Combined lab at NASA Ames



First laboratory to demonstrate simultaneous astrometry and high-contrast imaging. (2×10^{-8} contrast @ 2.0 to $3.5 \lambda/D$)

- Demonstrate real mission configuration
- Increase astrometry fidelity
- Coronagraph independent




Combined lab at NASA Ames



First laboratory to demonstrate simultaneous astrometry and high-contrast imaging. (TDEM 2013 Submitted)

- Demonstrate real mission configuration
- Increase astrometry fidelity
- Coronagraph independent (SP, HLC, Vortex, PIAA etc)





Combined lab at NASA Ames




- Preliminary results shows agreement with model

	Coronagraph field [arcsec]	Hexagon size a [μm]	Dot Size [μm]	1 st order location @ 632 nm	Spike width (350-900nm)
Angular location	$46 \lambda/D$	$42 \mu\text{m}$	$7 \mu\text{m}$	$254 \lambda/D$	$142 - 366 \lambda/D$
Focal plane	1 mm			5.1mm	3.1 - 7.94mm





Combined lab at NASA Ames



- Preliminary results shows agreement with model



	Coronagraph field [arcsec]	Hexagon size a [μm]	Dot Size [μm]	1 st order location @ 632 nm	Spike width (350-900nm)
Angular location	$46 \lambda/D$	$42 \mu\text{m}$	$7 \mu\text{m}$	$254 \lambda/D$	$142 - 366 \lambda/D$
Focal plane	1 mm			5.1mm	3.1 - 7.94mm



On sky results




- Carbon fiber pupil mask to test concept on sky.
- 1mas accuracy in a 1m telescope (No AO)
- Use of closest diffraction spike (reducing astrometric baseline) lowers systematic component by 2x



WFIRST



- **0.1 μ s imaging Astrometry**
- Simultaneous direct imaging
- Wide field telescope 0.3 $^{\circ}$
- Large aperture 2.4m
- No additional instrument is required, only a coating modification.




Astrometry accuracy: $\sim 0.1\mu$ s
Guyon et al., 2012

For a $D=2.4\text{m}$ and $\text{FoV}=0.3\text{ deg}^2$
0.098 μ s (Vis)
0.064 μ s (H)

Assuming:

2 Day integration
Galactic pole pointing
Telescope roll
Diffraction limited PSF




AFTA Telescope: Courtesy NASA

Outer Barrel Assembly



Optical Telescope Assembly





Support Structure



WFIRST



- The diffractive pupil is compatible with a general astrophysics mission.
- A 1% Diffractive pupil covers $\sim 1/5^{\text{th}}$ of the area of the spiders.
- Distortion calibration still be used **without** a bright star!





FoV %	Flux _{spike} /Flux _{Zodi} ratio vs star magnitudes		
	mV=6	mV=3.7	mV=-1.46
99%	$< 5 \times 10^{-2}$	$< 4 \times 10^{-1}$	$< 4.5 \times 10^1$
95%	$< 1.1 \times 10^{-3}$	$< 8.8 \times 10^{-3}$	< 1
90%	$< 3.6 \times 10^{-4}$	$< 3.0 \times 10^{-3}$	$< 3.4 \times 10^{-1}$



WFIRST

- Combination of PASS and GAIA can deliver $<10\mu\text{s}$ astrometry. (*From D. Spergel - A. Reiss*).



Still need to fit detector orientation positions and orientations:

-6 Parameters x 18


Detectors.

-Diffraction from spiders can help.


-Spiders+Secondary telemetry needed.

-*WFIRST Preparatory Science (WPS)*

Explorer Class (EXO-S, EXO-C?)



- 1-m Class wide-field explorer mission would allow exo-earth astrometric measurements.



- SSS Independent, works with
-Internal coronagraphs (i.e. EXO-C)
-and Star shade (i.e. EXO-S)
- Kepler-sized $D=1\text{m}$ and $\text{FoV}=1\text{ deg}^2$
 $0.17\mu\text{as}$ (Vis)

Expected Single Measurement Astrometric Accuracy as a Function of Telescope Diameter and Field of View (Galactic Pole Pointing, 2 Day Integration)

Tel. Diam.	Field of View			
	0.03 deg ² ($\phi = 0.2\text{ deg}$)	0.1 deg ² ($\phi = 0.36\text{ deg}$)	0.3 deg ² ($\phi = 0.6\text{ deg}$)	1 deg ² ($\phi = 1.13\text{ deg}$)
1.0 m	0.99 μas	0.54 μas	0.31 μas	0.17 μas
1.4 m	0.62 μas	0.34 μas	0.20 μas	0.11 μas
2.0 m	0.38 μas	0.21 μas	0.12 μas	0.066 μas
2.8 m	0.24 μas	0.13 μas	0.076 μas	0.041 μas
4.0 m	0.15 μas	0.081 μas	0.047 μas	0.026 μas
5.7 m	0.092 μas	0.050 μas	0.029 μas	0.016 μas
8.0 m	0.059 μas	0.032 μas	0.019 μas	0.010 μas

Proposed Astrometry SAG



- Goals:
 - SCIENCE: Identify astrometry niches.
 - COMMUNITY: Gather, individual or institutions that are currently engaged in astrometry research in the U.S.
 - TECHNOLOGY: Identify the most promising astrometry technology.
 - MISSION OPORTUNITIES: Assess cost-benefit of adding astrometry to planned missions.
 - ROAD MAP: Propose long-term roadmap for astrometry to maximize science return.